

Contributions of MS metabolomics to gilthead sea bream (*Sparus aurata*) nutrition.
Serum fingerprinting of fish fed low fish meal and fish oil diets

Rubén Gil-Solsona¹, Josep Alvar Calduch-Giner², Jaime Nácher-Mestre^{1,2,3}, Leticia
Lacalle-Bergeron¹, Juan Vicente Sancho¹, Félix Hernández¹, Jaume Pérez-Sánchez²

¹Research Institute for Pesticides and Water (IUPA). Avda. Sos Baynat, s/n. University
Jaume I, 12071 Castellón, Spain.

²Institute of Aquaculture Torre de la Sal (IATS, CSIC), 12595 Ribera de Cabanes,
Castellón, Spain.

³University Center EDEM, Muelle de la Aduana s/n, 46024 Valencia. Spain

Corresponding author:

Jaume Pérez-Sánchez

Email: jaime.perez.sanchez@csic.es

Phone: +34 964319500 ext 233

ABSTRACT

The aim of this study was to evaluate the impact of fish meal (FM) and fish oil (FO) replacement by plant proteins and oils in the serum metabolome of two-year old gilthead sea bream (*Sparus aurata*) fed from early life stages with control and experimental diets. Randomly selected fish were overnight sampled and clotted serum was used for metabolomics fingerprinting by means of ultra-high performance liquid chromatography coupled to quadrupole time-of-flight mass spectrometry. More than 12,500 different m/z ions were detected, and Partial Least Squares-Discriminant analysis separated fish fed control and plant-based diets, with a 71% of variance explained and 44% of variance predicted by the two first components. After variable importance in projection (VIP) and Benjamini-Hochberg test correction filtering, 50 endogenous compounds were elucidated as highly discriminant features of dietary treatment. Most of them were lipid-related compounds and reflected the different fatty acid composition of dietary oils, whereas changes in N-acyl taurines, cytidine and nucleoside related compounds would indicate changes in tissue repair and DNA degradation processes. Untargeted analysis also identified some exogenous compounds as markers of marine and vegetable raw materials. In the case of hercynine (antioxidant fungi and mycobacteria product), this was exemplified by a close lineal association between circulating and feed levels. Targeted approaches were focused on vitamins and a clear reduction of B₁₂, indirectly assessed via methylmalonic acid levels, was found in fish fed vegetable diets. Conversely, serum riboflavin (B₂) and pantothenic acid (B₅) levels were consistently increased, which highlighted the close link between nutrition and gut microbiota.

Keywords: Fish nutrition; liquid chromatography; mass spectrometry; metabolomics; vitamins; microbiota; plant-based diets.

1. Introduction

Current stagnation of fish meal (FM) and fish oil (FO) production from wild fisheries limits further growth of aquaculture (Tacon and Metian, 2015). The most obvious alternatives are the plant ingredients, which are a common practice in salmonids and marine fish to reduce the reliance of European aquaculture on marine fishery resources. Major progress in this way has been achieved within AQUAMAX, and ARRAINA EU projects and data on key performance indicators clearly indicate that alternative feeds with less than 7% marine ingredients support the maximum growth of gilthead sea bream (*Sparus aurata*) from early life stages to completion of sexual maturation (Benedito-Palos et al., 2016; Simó-Mirabet et al., 2018). It is also noteworthy that plant-based diets did not have a negative impact on the shelf life of gilthead bream, trout or carp as high quality foods (Grigorakis et al., 2018). Also, both in salmon and gilthead sea bream, no transfer from feeds to edible fillets was found for regulated mycotoxins, pesticides and persistent organic pollutants with the current plant-based diet formulations (Berntssen et al., 2005, 2010; Nacher-Mestre et al., 2009; 2015; Bell et al., 2012; Portolés et al., 2017). However, regardless of fish fatty acid (FA) biosynthetic capabilities, the use of plant-based diets is associated with a reduced content of n-3 long-chain poly-unsaturated FAs (PUFA) in the meat of farmed fish (Benedito-Palos et al., 2009; Liland et al., 2013; Ballester-Lozano et al., 2016; Turchini et al., 2018).

Other drawback effects of plant-based diets in marine farmed fish are related to changes in fish health and stress resilience (Montero and Izquierdo, 2010). Certainly, the magnitude and persistence of high plasma cortisol levels after crowding exposure is increased in juveniles of gilthead sea bream fed vegetable oils (Ganga et al., 2011), although a lower risk of oxidative stress in these challenged fish is also inferred (Pérez-Sánchez et al., 2013b). However, below the threshold level for the theoretical requirements in essential FAs, high inclusion levels of vegetable oils allow a faster disease progression in juveniles of gilthead sea bream challenged with the intestinal parasite *Enteromyxum leei* (Estensoro et al., 2011; Caldach-Giner et al., 2012). A possible cause are the nutritionally-mediated changes on the intestinal profile of mucins, mucosal immunoglobulins (*IgT*) and other immune-relevant genes of either diagnostic or predictive value (Caldach-Giner et al., 2012; Pérez-Sánchez et al., 2013c; Piazzon et al., 2016), which revealed a pro-inflammatory condition affecting also the integrity of

the intestinal barrier (Estensoro et al., 2016) and the composition of gut microbiota and intestinal mucus proteome (Piazzon et al., 2017). From these studies, however, it was also conclusive that most of these disturbing effects are reversed by the supplementation of plant-based diets with sodium butyrate, resulting in improved diseases outcomes in fish challenged with *E. ictaluri* and the bacteria *Photobacterium damsela* subsp. *piscicida* (Piazzon et al., 2017). Experimental evidence also indicates that diets enriched with medium-chain fatty acid salts (sodium heptanoate, sodium dodecanoate) have a positive impact on feed intake and energy metabolism of juvenile fish reared under sub-optimal conditions (Simó-Mirabet et al., 2017; Martos-Sitcha et al., 2018), although possible mechanisms still await full elucidation.

Very often, the application of targeted analyses is the prevailing strategy for qualitative and quantitative detection of different biomarkers. However, this strategy restricts the possibilities to detect other unpredictable effects that could result directly or indirectly from the changes in diet composition. This limitation has encouraged the development and application of new and powerful analytical approaches to face the complexity of this problem and to improve the chance to detect unanticipated effects. Currently a promising new “omic” approach is metabolomics, which aims to use profiles of low-molecular weight metabolic entities (usually < 1,000 Da) to identify biomarkers indicative of specific conditions and particular metabolic pathways. The novelty of this approach in aquaculture research is highlighted in the review article of Alfaro and Young (2018). In particular, nuclear magnetic resonance (NMR)-based lipid fingerprinting allows to precisely classify wild and farmed gilthead sea bream based on their muscle lipid composition (Melis et al., 2014). In another gilthead sea bream study, Robles et al. (2013) measured over 80 metabolites from fish intestine samples using a high-performance liquid chromatography-mass spectroscopy (HPLC–MS) platform. Although both analytical platforms rely on wide-untargeted approaches, MS allows retrospective analysis and a higher sensitivity and resolution power (Castro-Puyana and Herrero, 2013). Indeed, we have detected more than 15,000 *m/z* ions in the serum of gilthead sea bream by means of ultra-high performance liquid chromatography (UHPLC) and high resolution MS (HRMS) (Gil-Solsona et al., 2017). The same platform has been used in the present study to analyse fish from the eight-months feeding trial of Benedito-Palos et al. (2016). That study was prolonged, and herein data on wide- and targeted-serum metabolome were used to underline the effects of alternative feeds in two-year old fish fed experimental diets from early life stages.

2. Materials and methods

2.1. Reagents and chemicals

HPLC-grade methanol (MeOH), HPLC-supergradient acetonitrile (ACN), sodium hydroxide (> 99%), ammonium hydroxide (NH₄OH) and ammonium acetate (NH₄Ac) were obtained from Scharlab (Barcelona, Spain). HPLC-grade water was obtained from a Milli-Q water purification system (Millipore Ltd., Bedford, MA, USA). Leucine-enkephalin (mass-axis calibration), formic acid (mobile phase modifier), N,N-dimethyl L-histidine (reagent grade), methyl iodine (reagent grade) and tetrabutylammonium acetate (reagent grade) were purchased from Sigma-Aldrich (Saint Louis, MO, USA).

2.2. Diets

Four experimental diets were formulated and produced by BioMar (Brande, Denmark). All diets were isonitrogenous, isolipidic and isoenergetic and met all known nutritional requirements of gilthead sea bream. FM was included at 23% in the D1 (control) diet and at 3% in the other three experimental diets (D2, D3 and D4). Fish hydrolysate (CPSP) was added at 2% in all diets. Added oil was either FO (D1 diet) or a blend of vegetable oils (1:1 ratio of rapeseed oil: palm oil), replacing 58% (D2 diet) and 84% (D3 and D4 diets) FO. A commercial butyrate preparation (BP-70[®], NOREL) was added to the D4 diet at 0.4%. All diets contained histidine (0.14%), antioxidants (0.045%) and a mineral-vitamin mix (0.5%). Lysine, methionine, choline, lecithin and monocalcium phosphate were balanced in D2, D3 and D4 diets to the values of the control diet (Table 1).

2.3. Animal care and sampling

Juvenile fish (15 g initial average body weight) of Atlantic origin (Ferme Marine de Douhet, Ile d'Oléron, France) were fed control and experimental diets in the indoor experimental facilities of the Institute of Aquaculture Torre de la Sal (IATS-CSIC, Spain). Fish were allocated in 2,500 L tanks in triplicated groups (150 fish/tank), and each one was fed one of the experimental diets for 16 months from May 2013 to August

2014. The number of fish per tank was progressively reduced by periodical samplings, maintaining the rearing density below 15 kg/m³. Oxygen content of outlet water remained higher than 75% saturation and day-length and water temperature followed natural changes at IATS-CSIC latitude (40° 5'N; 0° 10'E). At time of sampling, actively fed fish (3-4 fish per tank to achieve 10 fish per diet) were sampled following overnight fasting for blood and tissue collection. Liver and visceral adipose tissue were extracted and weighed. Blood was taken from caudal vessels with vacutainer tubes with a clot activator, allowed to clot for 30 min at room temperature, and then centrifuged at 1,300 g for 10 min. The obtained samples were stored at -20°C until analysis.

All procedures were approved by the IATS Ethics and Animal Welfare Committee according to national (Royal Decree RD53/2013) and EU legislation (2010/63/EU) on the handling of animals for experiments.

2.4. UHPLC-HRMS

The analytical procedure was similar to that described elsewhere by Gil-Solsona et al. (2017). Briefly, serum samples were deproteinized with ACN and one supernatant aliquot was used for hydrophilic interaction liquid chromatography (HILIC). Another aliquot was evaporated to dryness and re-dissolved in MeOH 10% for reversed phase (RP) analysis. Quality control (QC) samples were prepared by pooling 50 µL of each sample extract. Extracts (10 µL) were injected in HILIC and RP in both positive and negative ionization modes (0.7 kV and 1.5 kV capillary voltages, respectively) in a hybrid quadrupole time-of-flight mass spectrometer (Xevo G2 QTOF, Waters, Manchester, UK) with a cone voltage of 25 V, using nitrogen as both desolvation and nebulizing gas.

The HILIC separation was performed using a mix of ACN:H₂O (95:5, v/v) as weak mobile phase (A) and H₂O as strong mobile phase (B) both in 0.01% formic acid (HCOOH) and 10 mM NH₄Ac. The percentage of B was changed as follows: 0 min, 2%; 1.5 min, 2%; 2.5 min, 15%; 6 min, 50%; 7.5 min, 75%; and finally at 7.51 min, 2%, with a total run time of 10 min, for both ESI+ and ESI-. For RP separation, the weak mobile phase (A) was H₂O with 0.01% HCOOH and the strong mobile phase (B) was MeOH with 0.01% HCOOH. The B percentage was changed from 10% at 0 min, to 90% at 14 min, 90% at 16 min and 10% at 16.01 min, with a total run time of 18 min for both ESI+ and ESI-. In order to obtain a better resolution among isomers of free FAs

and phospholipids, aliquots of RP samples were fortified at 50 mM with tetrabutylammonium acetate (TBA) and injected with the following gradient: A: H₂O 0.01% HCOOH, B: MeOH 0.01% HCOOH; The percentage of B was maintained at 70% during the first 5 min and changed from 70% at 5 min, to 80% at 8 min, 85% at 12 min, 95% at 15 min, 100% at 22 min and 70% again at 22.01 min with a total run time of 24 min for both ESI+ and ESI-.

2.5. Untargeted Data Processing

LC-MS data were processed using XCMS R package (<https://xcmsonline.scripps.edu/>) with *Centwave* algorithm for peak picking (peak width from 5 to 20 s, S/N ratio higher than 10 and mass tolerance of 15 ppm), followed by retention time alignment, peak area normalization (mean centering), log 2 applying (to avoid heteroscedasticity) and Pareto scaling. For elucidation purposes, fragmentation spectra of features of interest were compared with reference spectra databases (METLIN, <http://metlin.scripps.edu>; Human Metabolome DataBase, <http://www.hmdb.ca>; MassBank, <http://www.massbank.eu>). For unassigned metabolites, *in silico* fragmentation software (MetFrag, <http://msbi.ipb-halle.de/MetFrag>), with subsequent searches through Chemspider (<http://www.chemspider.com>) and PubChem (<https://pubchem.ncbi.nlm.nih.gov>) chemical databases, was employed.

2.6. Targeted analysis

The retrospective analysis of data acquired in MS^E mode served for the refined search of additional relevant compounds. This procedure consisted in the search of the *m/z* ratio (parent ions) of the metabolites of interest in the LE function, as well as product ions obtained from MS/MS spectrum online databases (METLIN and Human Metabolome DataBase) in the HE function. In the case of vitamins and related-compounds, fat-soluble vitamins were not directly analysable in serum, and their related metabolites were analysed as retinol phosphate for vitamin A, 25-hydroxyvitamin D₃ for vitamin D₃, α -Carboxyethylhydroxychroman for vitamin E and menaquinone for vitamin K₂ (Tai et al., 2010; Lebold et al., 2012; Karl et al., 2014). Water-soluble

vitamins were directly analysed (B₁, B₂, B₅, B₆, B₇ and C) with the exception of B₁₂, which was indirectly assayed as methylmalonic acid (MMA) (Lewerin et al., 2003).

Targeted analysis was also applied for hercynine, a betaine compound synthesized by fungi and mycobacteria. This exogenous compound was analysed in feeds and serum samples, using a hercynine standard synthesized as described elsewhere (Khonde and Jardine, 2015). In the case of feed samples, the analytical protocol included a polar extraction procedure previously employed in our laboratory for animal by-products (Nácher-Mestre et al., 2016). Briefly, 2.5 g of feeds were extracted with 10 mL H₂O:ACN (20:80) 0.1% HCOOH, centrifuged and supernatant (5 mL) was passed through an OASIS WCX SPE cartridge previously cleaned with 6 mL MeOH and 6 mL of Milli-Q H₂O. Sample was loaded, cleaned with 6 mL of MeOH:H₂O (1:1) and finally eluted in 3 mL of 2% formic acid in methanol. The feeds samples were then lead to dryness and diluted in 200 µL of Milli-Q H₂O to continue with MS analysis.

2.7. Statistical analysis

Data on growth performance and targeted analysis were analysed by one-way ANOVA followed by the Student Newman–Keuls test ($P < 0.05$). After data preprocessing of untargeted metabolomics, multivariate analysis was performed to find discriminative features among groups by means of the EZ-Info software (Umetrics, Sweden). First, Principal Component Analysis (PCA) was used to ensure the absence of outliers and the correct classification of QCs after normalization. Then, all the four experimental groups were joined in a single file and Partial Least Squares-Discriminant Analysis (PLS-DA) was conducted to maximize the separation of dietary groups. The contribution of m/z features to the PLS-DA model was assessed by means of variable importance in projection (VIP) measurements. A VIP score > 1 was considered an adequate threshold to determine discriminant variables in the PLS-DA model (Wold et al., 2001; Li et al., 2012; Kieffer et al., 2016). Additionally, orthogonal PLS-DA (Wiklund et al., 2008) with a high threshold ($P [\text{corr}] > 0.7$) was carried out to highlight the most discriminant compounds. To end, differences in normalized peak areas of m/z features were analysed by One-way ANOVA followed by Benjamini-Hochberg multiple testing correction analysis (Benjamini and Hochberg, 1995).

3. Results and Discussion

3.1. Fish condition

In the previous study of Benedito-Palos et al. (2016), data on key performance indicators and gene expression of growth-related markers in liver and skeletal muscle highly supported the suitability of FM/FO replacement by plant ingredients. In agreement with this, when fish coming from this initial trial were randomly sampled for serum metabolomics fingerprinting, all fish showed a similar average body weight ranging between 577 and 612 g (Table 2). Likewise, hepatosomatic index (HSI) and mesenteric fat index (MSI) remained mostly within the normal range of variation for the class of fish size and season (Cruz-García et al., 2009; Benedito-Palos et al., 2010). This revealed a lack of impact of dietary treatment upon body fat storage or tissue lipid trafficking, which are now recognized as clear signs of essential FA deficiencies in gilthead sea bream (Pérez-Sánchez et al., 2013a; Ballester-Lozano et al., 2015). Despite this, integrative omics approaches combining transcriptomics, proteomics and microbiome analyses highlighted a pro-inflammatory phenotype, with changes in the integrity of the epithelial intestinal barrier and diseases outcomes when fish fed plant-based diets are challenged with bacteria and enteric parasites (Estensoro et al., 2016; Piazzon et al., 2016; 2017). Recently, it has also been proven that plasma levels of sex steroids and the male-female sex reversal through the life cycle of gilthead sea bream are differentially regulated in fish fed marine and vegetable diets (Simo-Mirabet et al., 2018). Nevertheless, sex steroids (testosterone, 11-ketotestosterone, 17 β -estradiol) cannot be considered a major discriminating factor in this study, since their plasmatic concentrations increase gradually through gametogenesis in concomitance with gonadal growth, decreasing abruptly thereafter. Accordingly, circulating sex steroids were almost undetectable in our experimental setup using fish sampled out of the reproductive period, which normally extends for gilthead sea bream from October to March in our latitude (Chaoui et al., 2006; Hadj-Taieb et al., 2013). In any case, our methodology allowed a wide-screening approach, and a total of 12,982 m/z features (ions) were obtained in all four acquisition modes (RP and HILIC in both ionization modes ESI+ and ESI-). These numbers are comparable to those previously reported for fed and fasted juveniles of gilthead sea bream, using the same UHPLC-HRMS platform (Gil-Solsona et al., 2017). Of course, not all features corresponded to a single compound, but the number of detectable ions (13,000-15,000) was high enough to have

a wide-representation of the serum fish metabolome. Indeed, the number of different compounds in animal biofluids is estimated to be more than 8,000 (Kałużna-Czaplińska et al., 2014), with around 4,500 in human blood according to the Human Metabolome DataBase (Wishart et al., 2013).

3.2. Untargeted fingerprinting: multivariate analysis

One-way ANOVA was suitable to detect a wide-range of changes in circulating metabolites with more than 5,000 differentially expressed ions when comparing control and extreme D3/D4 groups ($P < 0.05$), but these numbers were drastically reduced after filtering with Benjamini-Hochberg for false positive corrections (Fig. 1A). Thus, the number of ions with a different abundance ranged between 451 and 2,929 when comparisons were made between D1 and D2 fish; and D1 and D4 fish, respectively. However, only four individual features were different between groups D3 and D4, which was indicative that the source of variation when FM/FO diets were supplemented with butyrate was very low in comparison to that of the replacement of FM and FO with plant ingredients alone. This was also evidenced by multivariate PLS-DA analysis as many individuals of D3 and D4 groups overlapped in the score plot (Fig. 1B). This is the reason why data from fish fed D3 and D4 were pooled in the same group (D3/4) for subsequent PLS-DA analyses, where the 71% of variance and 44% of variance was explained or predicted, respectively, by the two first components. The maximum individual variability was achieved within D2 group, but importantly all fish of D1 and D3/D4 groups were correctly classified in the discriminant model. Thus, the maximum separation along both components was found for D1 and D3/4 fish that were distributed along the first (X-axis) and second (Y-axis) component, whereas the separation of D2 and D3/4 fish was only evidenced along the first component reflecting the changes in FO inclusion levels (6.5% D2 diet; 2.50% D3/4 diets). In contrast, the distribution along the second component would primarily reflect the reduced feed intake of FM and fish hydrolysates with inclusion levels of 25.0% in D1 diet and 5.0% in D2, D3 and D4 diets. However, it is noteworthy that the number of features with a $P[\text{corr}] > 0.95$ by Orthogonal PLS-DA was reduced to 39, whereas up to 850 ions were identified as clearly discriminant ions in 10-days fasted fish (Gil-Solsona et al., 2017). Therefore, the magnitude of changes induced in the present study by dietary treatment were markedly reduced in comparison to the fasting mediated effects, which suggests that most of them

primarily mirror differences in diet composition rather than functional metabolic dysfunctions associated to changes in diet composition. For this reason, a less restrictive $P[\text{corr}] > 0.7$ was used for the subsequent elucidation procedures.

3.3. Elucidation of untargeted differential compounds

A total of 55 representative compounds with statistically significant changes in abundance after correction for false positives and a VIP score > 1.3 were elucidated (Table 3). Most of them were compounds of lipid nature, such as phosphocholines (PC, 24), lysophosphocholines (LysoPC, 10), free FAs (8) and sphingolipids (2). Other compounds with a different abundance were elucidated as N-acyl-taurines (2), cytidine and cytosine nucleosides (4), cysteinolic acid, tauropine, trimethylamine N-oxide (TMAO), arsenobetaine and hercynine. Accordingly, most of these compounds are related to lipid metabolism and highly reflected the decreased unsaturation index of FAs of vegetable oils. Indeed, FO has an elevated content of n-3 LC-PUFAs, whereas vegetable oils are almost devoid of eicosapentaenoic acid (20:5n-3) and docosahexaenoic acid (22:6n-3), which cannot be synthesized at a high rate in marine fish from the C18 PUFA precursor, α -linolenic acid (18:3n-3) (Tocher, 2015). In consequence, previous studies in gilthead sea bream clearly indicate that the inclusion of vegetable oils in fish feeds reduced the content of LC-PUFAs and increased that of C18 PUFAs in liver, adipose tissue and muscle fillets, with a selective incorporation of unsaturated FAs in polar lipids (Izquierdo et al., 2005; Benedito-Palos et al., 2010; 2013) to preserve and maintain the function of cell membrane surfaces. Especially for fat fish species, most of these changes in the flesh FA composition are highly predictable by means of a dummy regression model (Ballester-Lozano et al., 2014; 2016). Less is known about the effects of diet composition on the FA composition of circulating lipids, although clinical studies evidence that they also reflect the changes in diet composition (Laidlaw and Holub, 2003; Lemaitre et al., 2003) as it was herein the case of circulating PCs, lysoPCs and free FAs. Moreover, the number and degree of these changes in comparison to control group D1 increase with the level of replacement in a dose-dependent manner.

Sphingolipids, as well as phospholipids, are essential components of all eukaryotic cell membranes with important roles in a variety of biological processes including cell division and cell-to-cell interactions (Hannun and Obeid, 2018). In their

simplest forms, sphingosine, phytosphingosine, and dihydrosphingosine serve as the backbones upon which further complexity is achieved. For example, phosphorylation of the C1 hydroxyl group yields the final breakdown products and/or the important signalling molecules sphingosine-1-phosphate, phytosphingosine-1-phosphate and dihydrosphingosine-1-phosphate, respectively (Gault and Obeid, 2010). In the present study, two sphingosine-related compounds were altered by dietary treatment and the abundance of (9-methyl-d19:3) sphingosine was markedly reduced (D2, 23% control fish; D3, 16% control) by the replacement of marine resources by plant ingredients. Conversely, (d14:2) sphingosine was markedly increased in D2 fish (878% with respect to D1 group) with intermediate values with the extreme diet formulation in D3/4 fish, which suggests that other factors than the simple inclusion level of plant ingredients have effects on sphingolipid metabolism, but we are still far to understand the physiological significance of this finding.

In recent years, a number of studies have demonstrated the essentiality of dietary taurine for many commercially relevant species, especially marine teleosts. Consequently, the removal of taurine-rich dietary ingredients such as FM can induce deficiencies with a wide range of symptoms, including reduced growth and survival, increased susceptibility to diseases and impaired larval developments as reviewed by Salze and Davis (2015). However, the paradigm that taurine is an essential nutrient is not nearly as clear in freshwater species and it is difficult to draw definitive conclusions, although the list of fish species for which taurine is required is increasing. In any case, taurine is well recognized as an essential nutrient in most carnivorous fish, and early studies in gilthead sea bream indicated that low levels of taurine in the pool of muscle free amino acids is associated with growth impairments in fish fed plant protein-based diets (Gómez-Requeni et al., 2004). The amides of long-chain FAs with taurine (N-acyl-taurines) are produced via oxidation of bile acid precursors in peroxisomes, and can function as cell signalling molecules with a wide range of biological activities (Hunt et al., 2012). N-acyl-taurines have been recently identified in liver and other rodent tissues, and genetic deletion or pharmacological blockage of the serine amidase FA amide hydrolase (FAAH) causes profound acceleration on wound healing in mouse skin, and repair associated responses in primary cultures of human keratinocytes and fibroblasts (Sasso et al., 2016). In the same study, immunofluorescence images of intact mouse skin show that FAAH co-localizes with cytokeratin 10 and filaggrin, two proteins that are expressed by epidermal supra-basal keratinocytes. In a previous study, we have

identified the cytokeratin 8 as a good marker of multiple aquaculture stressors (tank shaking, sounds, moving objects into water, water reverse flow and light flashes) in the skin mucus of gilthead sea bream (Pérez-Sánchez et al., 2017). The association of cytokeratins with N-acyl taurines has not been established in fish, but we found herein that the concentration of either N-heptadecenoyl-aurine or N-palmitoleoyl-aurine was progressively and consistently reduced with the combined replacement of FM and FO by plant ingredients. This finding opens new research issues in fish nutrition, which would be targeted to alleviate some of the drawback effects of plant-based diets upon the epithelial mucosae of gilthead sea bream, probably mediated by cell renewal or anti-inflammatory processes, as it has been reported for other bioactive compounds, such as butyrate which helps to restore and preserve the integrity and function in gilthead sea bream fed from early life stages with plant-based diets (Estensoro et al., 2016; Piazzon et al., 2017). Moreover, experimental evidence indicates that both butyrate and taurine are able to mitigate through different modes of action the intestinal anomalies of European sea bass fed with highly enriched soybean meal diets (Rimoldi et al., 2016).

Cytidine and nucleoside related compounds (cytosine, deoxycytidine, methylcytosine) were also clear discriminant factors in our experimental model, and their concentrations were consistently increased in fish fed plant-based diets. Intriguingly this was more evident in the group of fish fed D2 diet (200-734% control fish) than in the extreme D3/4 group (120-190% control fish). Since these compounds originate from dietary sources, from cellular excretion subsequent to RNA turnover, from cytosolic pools of nucleotides, or from degradation of nuclear DNA phagocytized by macrophages (Holstege et al., 1984), it is difficult to understand the physiological significance of these findings, although a major source of variation might be related to some kind of cellular DNA instability. Indeed, the highest difference amount control and experimental groups was reported for deoxycytidine and methylcytosine. Degradation of DNA produces deoxycytidine and chemotherapy sharply raises plasma deoxycytidine levels above pretreatment levels (Cohen et al., 1997). At the same time, methylation of cytosines is an important element of epigenetic regulation, and the increased circulating levels of methylcytosine can indicate not only a higher DNA degradation or instability, but also a hyper-methylation at the whole DNA or at specific gene sites. However, this notion needs to be confirmed by more specific assays, because vegetarian life styles are associated with hypo-methylation states (Geisel et al., 2005).

Unlike endogenous compounds, the origin and significance of exogenous compounds with a different abundance was easier to trace, being highly informative of the nature and origin of feed ingredients. Accordingly, the replacement of FM by plant ingredients was associated to a decrease of circulating cysteinolic acid, tauropine, TMAO or arsenobetaine. Cysteinolic acid is a non-protein amino acid similar to taurine, detected in gilthead sea bream and red sea bream (*Pagrus major*) as cholesterol-conjugate precursors in the synthesis of bile salts (Goto et al., 1996; Une et al., 1991). This amino acid is not synthesized by fish, but it can be easily incorporated in the food chain as some marine seaweed such as *Ulva* or *Enteromorpha* contain large amounts (Ito, 1963). Likewise, tauropine is an anaerobic end product found in several marine invertebrate phyla, but widely prevalent in marine molluscs (Venter et al., 2016). The same for TMAO, a compound found in animals, plants and fungi, but the concentration of TMAO in marine animals significantly exceeds that of other organisms (Yancey, 2005). Likewise, arsenobetaine is the arsenic analogue of the quaternary ammonium compound glycine betaine, and marine animals contain very high levels of this compound, non-toxic for human or animals (Molin et al., 2015; Stiboller et al., 2015). Its relative contribution of trophic transfer and biotransformation of arsenic derivatives in the arsenobetaine content in fish is still under debate (Caumette et al., 2012; Popowich et al., 2016), although from our results it was evident the direct relation between dietary FM and circulating arsenobetaine levels.

Another exogenous compound with a high discriminant value in our experimental model was hercynine. This is an intermediate compound in the synthesis of ergothioneine, a natural antioxidant that is only synthesized by non-yeast fungi, cyanobacteria and actinobacteria (Fahey, 2001; Pfeiffer et al., 2011). Therefore, its detectable presence in the serum of fish is indicative of feeding plant ingredients, although its circulating concentration did not parallel the replacement level, being the circulating concentration (arbitrary units) in D2 fish ($737 \pm 74\%$) too much higher than that of D3/4 ($182 \pm 15\%$) fish. However, when these values were plotted against the relative concentration of hercynine in the diet, a close linear association was found for this compound (Fig. 2). Therefore, with the advent of new formulations, hercynine is coming as good biomarker of raw material traceability, but also of proper feed storage and processing of plant-based diets with no fungi/mycobacteria growth.

3.4. Targeted vitamin analysis

Vitamins are essential micronutrients that are normally found as precursors of various enzyme reactions in all living cells. However, most of them cannot be synthesized by animals and they need to be obtained exogenously by means of diet fortification, although the use of vitamin-producing microorganisms represents a more natural and consumer-friendly alternative (Le Blanc et. al., 2013). In humans, it has been shown that members of the gut microbiota are able to synthesize vitamin K as well as most of the water-soluble B vitamins, such as biotin, cobalamin, folates, nicotinic acid, pantothenic acid, pyridoxine, riboflavin and thiamine (Hill, 1997). Unlike dietary vitamins, the predominant uptake of the microbially-produced vitamins occurs in the colon (Said and Mohammed, 2016). A similar specialization seems to exist along the digestive tract of fish, as evidenced the microarray gene expression profiling of several genes related to vitamin B₁₂ through the intestine of European sea bass (Calduch-Giner et al., 2016). Experimental evidence also indicates that replacement of FM by plant ingredients drives many changes in the micronutrient diet composition, with an important decrease in the content of some vitamins (NRC, 2011). In our experimental model, most of the theoretically mineral and vitamin requirements are met in excess by the diet (Table 1), but to assess the proper levels of circulating vitamins and vitamin-related compounds, a retrospective (targeted) analysis was conducted by means of the MS^E acquisition mode. This approach served to check deficiencies in specific compounds that could have been masked by the astringent Benjamini-Hochberg multiple testing correction in the untargeted approach. Hence, as shown in Table 4, the relative concentration of riboflavin (vitamin B₂) and pantothenic acid (vitamin B₅) were progressively and significantly increased with the replacement of marine sources by plant ingredients in D3/4 fish. Conversely, methylmalonic acid (MMA), used as a biomarker of vitamin B₁₂ deficiency in humans and rodents (Watanabe et al., 1991; Carmel, 2011), increased progressively and significantly with the replacement FM/FO by plant ingredients in fish fed D2 and D3/4 diets. The replacement of FM by plant proteins also decreased the concentration of vitamin B₁₂ in muscle and liver tissues of Atlantic cod (Hansen et al., 2007), being now well recognized the risk of vitamin B₁₂ deficiency in vegetarian humans (Stabler and Allen, 2004; Allen, 2009). Our targeted approach did not detect additional changes in vitamin condition, although vitamin B₇ is markedly reduced by short-term fasting in gilthead sea bream (Gil-Solsona et al., 2017).

482 All this reinforces the importance to define the core microbiota for a given feeding
483 regime and nutritional status, but studies in livestock animal and fish in particular are
484 still in an infancy state to fully understand the complexity of host and gut microbiota
485 interactions.

486

487

4. Conclusions

UHPLC-HRMS approach allowed us to identify a high number of m/z ions in the serum of farmed gilthead sea bream. This was the result of combined targeted and untargeted approaches, which identified a wide-range of endogenous and exogenous compounds with a high discriminant capacity as summarized in Fig. 3. Multivariate analyses highlighted a clear separation of fish fed the control and plant-based diets, and the distribution through X-axis and Y-axis evidenced the different effects related to FM or FO replacement by plant proteins and oils. Most of the changes reflected the different FA composition of dietary oils in fish growing at high rates without apparent signs of FA deficiencies. However, N-acyl taurines emerged as target compounds to alleviate some of the negative health effects of plant-based diets. Other metabolite changes (cytidine and nucleoside compounds) highlighted different nutritionally-mediated effects on DNA stability and perhaps methylation levels. Targeted vitamin analysis corroborated the risk of low levels of vitamin B₁₂ in fish fed plant-based diets, whereas other dietary or microbially-produced vitamins were not affected or increased (B₂, B₅). Lastly, the detection of different exogenous compounds served to trace the use of different raw materials in fish feeds, but also to eventually assess their proper processing and storage.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

Author contributions

J.V.S, F.H. and J.P.S conceived and designed the experiments. R.G.S, J.C.G., J.N.M., L.L.B. and J.P.S. performed the experiments. All authors have contributed to analysis of data and the final writing of the paper. All authors have read and approved the final manuscript.

Acknowledgments

The authors acknowledge the financial support of the Spanish MINECO (MI2-Fish, AGL2013- 48560) and Pla de Promoció de la Investigació de la Universitat Jaume

522 I (P1· 1B2015-59). Additional funding was obtained from Generalitat Valenciana, as
523 research groups of excellence (PROMETEO II/2014/023, PROMETEO II/2014/085,
524 Collaborative Research on Environment and Food-Safety, ISIC/2012/016). This work
525 has been developed in the framework of the Research Unit of Marine Ecotoxicology
526 (IATS-CSIC; IUPA-UJI).

References

- Alfaro, A.C., Young, T., 2018. Showcasing metabolomics research in aquaculture: a review. *Reviews in Aquaculture* 10, 135-152.
- Allen, L.H., 2009. How common is vitamin B12 deficiency? *Am. J. Clin. Nutr.* 89, S693-S696.
- Ballester-Lozano, G.F., Benedito-Palos, L., Estensoro, I., Sitjà-Bobadilla, A., Kaushik, S., Pérez-Sánchez J. 2015. Comprehensive biometric, biochemical and histopathological assessment of nutrient deficiencies in gilthead sea bream fed semi-purified diets. *British Journal of Nutrition* 114, 713-726.
- Ballester-Lozano, G.F., Benedito-Palos, L., Mingarro, M., Navarro, J.C., Pérez-Sánchez, J., 2016. Up-scaling validation of a dummy regression approach for predictive modelling the fillet fatty acid composition of cultured European sea bass (*Dicentrarchus labrax*). *Aquaculture Research* 47, 1067-1074.
- Ballester-Lozano, G.F., Benedito-Palos, L., Riaza, A., Navarro, J.C., Rosel, J., Pérez-Sánchez, J., 2014. Dummy regression analysis for modelling the nutritionally tailored fillet fatty acid composition of turbot and sole using gilthead sea bream as a reference subgroup category. *Aquaculture Nutrition* 20, 421-430.
- Bell, J.G., Dick, J.R., Strachan, F., Guy, D.R., Berntssen, M.H.G., Sprague, M., 2012. Complete replacement of fish oil with a blend of vegetable oils affects dioxin, dioxin-like polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in 3 Atlantic salmon (*Salmo salar*) families differing in flesh adiposity. *Aquaculture* 324-325, 118-126.
- Benedito-Palos, L., Ballester-Lozano, G.F., Simó, P., Karalazos, V., Ortiz, A., Calduch-Giner, J.A., Pérez-Sánchez, J., 2016. Lasting effects of butyrate and low FM/FO diets on growth performance, blood haematology/biochemistry and molecular growth-related markers in gilthead sea bream (*Sparus aurata*). *Aquaculture* 454, 8-18.
- Benedito-Palos, L., Calduch-Giner, J.A., Ballester-Lozano, G.F., Pérez-Sánchez, J., 2013. Effect of ration size on fillet fatty acid composition, phospholipid allostasis and mRNA expression patterns of lipid regulatory genes in gilthead sea bream (*Sparus aurata*). *British Journal of Nutrition* 109, 1175-1187.
- Benedito-Palos, L., Navarro, J.C., Bermejo-Nogales, A., Saera-Vila, A., Kaushik, S., Pérez-Sánchez, J., 2009. The time course of fish oil wash-out follows a simple dilution model in gilthead sea bream (*Sparus aurata* L.) fed graded levels of vegetable oils. *Aquaculture* 288, 98-105.
- Benedito-Palos, L., Navarro, J.C., Kaushik, S., Pérez-Sánchez, J., 2010. Tissue-specific robustness of fatty acid signatures in cultured gilthead sea bream (*Sparus aurata* L.) fed practical diets with a combined high replacement of fish meal and fish oil. *Journal of Animal Science* 88, 1759-1770.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B* 57, 289-300.
- Berntssen, M.H.G., Julshamn, K., Lundebye, A.K., 2010. Chemical contaminants in aquafeeds and Atlantic salmon (*Salmo salar*) following the use of traditional-versus alternative feed ingredients. *Chemosphere* 78, 637-646.

572 Berntssen, M.H.G., Lundebye, A.K., Torstensen, B.E., 2005. Reducing the levels of
573 dioxins and dioxin-like PCBs in farmed Atlantic salmon by substitution of fish oil
574 with vegetable oil in the feed. *Aquaculture Nutrition* 11, 219-231.

575 Calduch-Giner, J.A., Sitjà-Bobadilla, A., Davey, G.C., Cairns, M.T., Kaushik, S., Pérez-
576 Sánchez, J., 2012. Dietary vegetable oils do not alter the intestine transcriptome of
577 gilthead sea bream (*Sparus aurata*), but modulate the transcriptomic response to
578 infection with *Enteromyxum leei*. *BMC Genomics* 470, 13.

579 Calduch-Giner, J.A., Sitjà-Bobadilla, A., Pérez-Sánchez, J., 2016. Gene expression
580 profiling reveals functional specialization along the intestinal tract of a
581 carnivorous teleostean fish (*Dicentrarchus labrax*). *Frontiers in Physiology* 7,
582 359.

583 Carmel, R., 2011. Biomarkers of cobalamin (vitamin B-12) status in the epidemiologic
584 setting: a critical overview of context, applications, and performance
585 characteristics of cobalamin, methylmalonic acid, and holotranscobalamin II. *Am.*
586 *J. Clin. Nutr.* 94, 348S-358S..

587 Castro-Puyana, M., Herrero, M., 2013. Metabolomics approaches based on mass
588 spectrometry for food safety, quality and traceability, *Trends in Analytical*
589 *Chemistry* 52, 74-87.

590 Caumette, G., Koch, I., Reimer, K.J., 2012. Arsenobetaine formation in plankton: a
591 review of studies at the base of the aquatic food chain. *J. Environ. Monitor.* 14,
592 2841-2853.

593 Chaoui, L., Kara, M.H., Faure, E., Quignard, J.P., 2006. Growth and reproduction of the
594 gilthead seabream *Sparus aurata* in Mellah lagoon (north-eastern Algeria).
595 *Scientia Marina* 70, 545-552.

596 Cohen, J.D., Strock, D.J., Teik, J.E., Katz, T.B., Marcel, P.D., 1997. Deoxycytidine in
597 human plasma: Potential for protecting leukemic cells during chemotherapy.
598 *Cancer Lett.* 116, 167-175.

599 Cruz-García, L., Saera-Vila, A., Navarro, I., Calduch-Giner, J., Pérez-Sánchez, J., 2009.
600 Targets for the TNF alpha-induced lipolysis in gilthead sea bream (*Sparus aurata*
601 L.) adipocytes isolated from lean and fat juveniles fish. *J. Exp. Biol.* 212, 2254-
602 2260.

603 Estensoro, I., Ballester-Lozano, G.F., Benedito-Palos, L., Grammes, F., Martos-Sitcha,
604 J.A., Mydland, L.-T., Calduch-Giner, J.A., Fuentes, J., Karalazos, V., Ortiz, A.,
605 Øverland, M., Sitjà-Bobadilla, A., Pérez-Sánchez, J., 2016. Dietary butyrate helps
606 to restore the intestinal status of a marine teleost (*Sparus aurata*) fed extreme diets
607 low in fish meal and fish oil. *PLoS ONE* 11, e0166564.

608 Estensoro, I., Benedito-Palos, L., Palenzuela, O., Kaushik, S., Sitjà-Bobadilla, A.,
609 Pérez-Sánchez, J., 2011. The nutritional background of the host alters the disease
610 outcome in a fish-myxosporean system. *Veterinary Parasitology* 175, 141-150.

611 Fahey, R.C., 2001. Novel thiols of prokaryotes. *Annu. Rev. Microbiol.* 51, 333-356.

612 Ganga, R., Bell, J.G., Montero, D., Atalah, E., Vraskou, Y., Tort, L., Fernández, A.,
613 Izquierdo, M.S., 2011. Adrenocorticotrophic hormone-stimulated cortisol release
614 by the head kidney inter-renal tissue from sea bream (*Sparus aurata*) fed with
615 linseed oil and soybean oil. *British Journal of Nutrition* 105, 238-247.

616 Gault, C.R., Obeid, L.M., 2011. Still benched on its way to the bedside: sphingosine
617 kinase 1 as an emerging target in cancer chemotherapy. *Critical Reviews in*
618 *Biochemistry and Molecular Biology* 46, 342-351.

619 Geisel, J., Schorr, H., Bodis, M., Isber, S., Hübner, U., Knapp, J.P., Obeid, R.,
620 Herrmann, W., 2005. The vegetarian lifestyle and DNA methylation. *Clin. Chem.*
621 *Lab. Med.* 43, 1164-1169.

622 Gil-Solsona, R., Nacher-Mestre, J., Lacalle-Bergeron, L., Sancho, J.V., Calduch-Giner,
623 J.A., Hernández, F., Pérez-Sánchez, J., 2017. Untargeted metabolomics approach
624 for unraveling robust biomarkers of nutritional status in fasted gilthead sea bream
625 (*Sparus aurata*). *PeerJ* 5, e2920.

626 Gómez-Requeni, P., Mingarro, M., Calduch-Giner, J., Medale, F., Martín, S.A.M.,
627 Houlihan, D.F., Kaushik, S., Pérez-Sánchez, J., 2004. Protein growth
628 performance, amino acid utilisation and somatotrophic axis responsiveness to fish
629 meal replacement by plant protein sources in gilthead sea bream (*Sparus aurata*).
630 *Aquaculture* 232, 493-510.

631 Goto, T., Ui, T., Une, M., Kuramoto, T., Kihira, K., Hoshita, T., 1996. Bile salt
632 composition and distribution of the D-cysteinolic acid conjugated bile salts in fish.
633 *Fish. Sci.* 62, 606-609.

634 Grigorakis, K., Dimitra, K., Corraze, G., Pérez-Sánchez, J., Adorjan, A., Zsuzsanna,
635 J.S., 2018. Impact of diets containing plant raw materials as fish meal and fish oil
636 replacement on rainbow trout (*Oncorhynchus mykiss*), gilthead sea bream (*Sparus*
637 *aurata*), and common carp (*Cyprinus carpio*) freshness. *Journal of Food Quality*
638 2018, 1717465.

639 Hadj-Taieb, A., Ghorbel, M., Hadj-Hamida, N.B., Jarboui O. Sex ratio, reproduction,
640 and growth of the gilthead sea bream, *Sparus aurata* (Pisces: Sparidae), in the
641 Gulf of Gabes, Tunisia. *Ciencias Marinas* 39, 101-112.

642 Hannun, Y.A., Obeid, L.M., 2018. Sphingolipids and their metabolism in physiology
643 and disease. *Nature Reviews Molecular Cell Biology* 19, 175-191.

644 Hansen, A.C., Rosenlund, G., Karlsen, Ø., Koppe, W., Hemre, G.I., 2007. Total
645 replacement of fish meal with plant proteins in diets for Atlantic cod (*Gadus*
646 *morhua* L.). I. Effects on growth and protein retention. *Aquaculture* 272, 599-611.

647 Hill, M.J., 1997. Intestinal flora and endogenous vitamin synthesis. *Eur. J. Cancer Prev.*
648 6 (Suppl 1), S43-S45.

649 Holstege, A., Manglitz, D., Gerok, W., 1984. Depletion of blood plasma cytidine due to
650 increased hepatocellular salvage in d-galactosamine-treated rats. *Eur. J. Biochem.*
651 141, 339-344.

652 Hunt, M.C., Siponen, M.I., Alexson, S.E.H., 2012. The emerging role of acyl-CoA
653 thioesterases and acyltransferases in regulating peroxisomal lipid metabolism.
654 *Biochimica et Biophysica Acta* 1822, 1397-1410.

655 Ito, K., 1963. Distribution of d-cysteinolic acid in marine algae. *Bull. Jap. Soc. Scient.*
656 *Fish* 29, 771-775.

657 Izquierdo, M.S., Montero, D., Robaina, L., Caballero, R., Rosenlund, G., Ginés, R.,
658 2005. Alterations in fillet fatty acid profile and flesh quality in gilthead seabream
659 (*Sparus aurata*) fed vegetable oils for a long term period. *Recovery of fatty acid*

660 profiles by fish oil feeding. *Aquaculture* 250, 431-444.

661 Kałużna-Czaplińska, J., Jóźwik, J., Żurawicz, E., 2014. Analytical methods used in
662 autism spectrum disorders. *Trends in Analytical Chemistry* 62, 20-27.

663 Karl, J.P., Fu, X., Dolnikowski, G.G., Saltzman, E., Booth, S.L., 2014. Quantification of
664 phylloquinone and menaquinones in feces, serum, and food by high-performance
665 liquid chromatography-mass spectrometry. *Journal of Chromatography B* 963,
666 128-133.

667 Khonde, P.L., Jardine, A., 2015. Improved synthesis of the super antioxidant,
668 ergothioneine, and its biosynthetic pathway intermediates. *Organic &
669 Biomolecular Chemistry* 13, 1415-1419.

670 Kieffer, D.A., Piccolo, B.D., Vaziri, N.D., Liu, S., Lau, W.L., Khazaeli, M.,
671 Nazertehrani, S., Moore, M.E., Marco, M.L., Martin, R.J., Adams, S.H., 2016.
672 Resistant starch alters gut microbiome and metabolomic profiles concurrent with
673 amelioration of chronic kidney disease in rats. *Am. J. Physiol. Renal. Physiol.*
674 310, F857-F871.

675 Laidlaw, M., Holub, B.J., 2003. Effects of supplementation with fish oil-derived n-3
676 fatty acids and gamma-linolenic acid on circulating plasma lipids and fatty acid
677 profiles in women. *Am. J. Clin. Nutr.* 77, 37-42.

678 LeBlanc, J.G., Milani, C., de Giori, G.S., Sesma, F., van Sinderen, D., Ventura, M.,
679 2013. Bacteria as vitamin suppliers to their host: A gut microbiota perspective.
680 *Curr. Opin. Biotechnol.* 24, 160-168.

681 Lebold, K.M., Ang, A., Traber, M.G., Arab, L., 2012. Urinary α -carboxyethyl
682 hydroxychroman can be used as a predictor of α -tocopherol adequacy, as
683 demonstrated in the Energetics Study. *The American Journal of Clinical Nutrition*
684 96, 801-809.

685 Lemaitre, R.N., King, I.B., Mozaffarian, D., Kuller, L.H., Tracy, R.P., Siscovick, D.S.,
686 2003. n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal
687 myocardial infarction in older adults: the Cardiovascular Health Study. *Am. J.*
688 *Clin. Nutr.* 77, 319-325.

689 Lewerin, C., Nilsson-Ehle, H., Matousek, M., Lindstedt, G., Steen, B., 2003. Reduction
690 of plasma homocysteine and serum methylmalonate concentrations in apparently
691 healthy elderly subjects after treatment with folic acid, vitamin B₁₂ and vitamin
692 B₆: a randomised trial. *European Journal of Clinical Nutrition* 57, 1426-1436.

693 Li, H., Ma, M.L., Luo, S., Zhang, R.M., Han, P., Hu, W., 2012. Metabolic responses to
694 ethanol in *Saccharomyces cerevisiae* using a gas chromatography tandem mass
695 spectrometry-based metabolomics approach. *Int. J. Biochem. Cell. Biol.* 44, 1087-
696 1096.

697 Liland, N.S., Rosenlund, G., Berntssen, M.H.G., Brattelid, T., Madsen, L., Torstensen,
698 B.E., 2013. Net production of atlantic salmon (FIFO, fish in fish out < 1) with
699 dietary plant proteins and vegetable oils. *Aquaculture Nutrition* 19, 289-300.

700 Martos-Sitcha, J.A., Simó-Mirabet, P., Piazzon, M.C., de las Heras, V., Calduch-Giner,
701 J.A., Puyalto, M., Tinsley, J., Makol, A., Sitjà-Bobadilla, A., Pérez Sánchez, J.,
702 2018. Dietary sodium heptanoate helps to improve feed efficiency, growth
703 hormone status and swimming performance in gilthead sea bream (*Sparus*
704 *aurata*). *Aquaculture Nutrition* (in press).

705 Melis, R., Sanna, R., Braca, A., Bonaglini, E., Cappuccinelli, R., Slawski, H., Roggio,
706 T., Uzzau, S., Anedda, R., 2017. Molecular details on gilthead sea bream (*Sparus*
707 *aurata*) sensitivity to low water temperatures from ¹H NMR metabolomics.
708 *Comparative Biochemistry & Physiology Part A Molecular & Integrative*
709 *Physiology* 204, 129-136.

710 Molin, M., Ulven, S.M., Meltzer, H.M., Alexander, J., 2015. Arsenic in the human food
711 chain, biotransformation and toxicology - review focusing on seafood arsenic. *J.*
712 *Trace Elem. Med. Biol.* 31, 249-259.

713 Montero, D., Izquierdo, M.S., 2010. Welfare and health of fish fed vegetable oils as
714 alternative lipid sources to fish oil, in: Turchini, G., Ng, W., Tocher, D. (Eds),
715 *Fish oil replacement and alternative lipid sources in aquaculture feeds*. CRC
716 Press, Cambridge, pp. 439-485.

717 Náchter-Mestre, J., Ibáñez, M., Serrano, R., Boix, C., Bijlsma, L., Lunestad, B.T.,
718 Hannisdal, R., Alm, M., Hernández, F., Berntssen, M.H.G., 2016. Investigation of
719 pharmaceuticals in processed animal by-products by liquid chromatography
720 coupled to high-resolution mass spectrometry. *Chemosphere* 154, 231-239.

721 Náchter-Mestre, J., Serrano, R., Beltrán, E., Pérez-Sánchez, J., Silva, J., Karalazos, V.,
722 Hernández, F., Berntssen, M.H.G., 2015. Occurrence and transfer of mycotoxins
723 in gilthead sea bream and Atlantic salmon by use of novel alternative feed
724 ingredients. *Chemosphere* 128, 314-320.

725 Náchter-Mestre, J., Serrano, R., Benedito-Palos, L., Navarro, J.C., López, F.J., Pérez-
726 Sánchez, J., 2009. Effects of fish oil replacement and re-feeding on the
727 bioaccumulation of organochlorine compounds in gilthead sea bream (*Sparus*
728 *aurata* L.) of market size. *Chemosphere* 76, 811-817.

729 NRC, National Research Council, 2011. *Nutrient Requirement of Fish and Shellfish*,
730 National Academy Press, Washington.

731 Pérez-Sánchez, J., Benedito-Palos, L., Ballester-Lozano, G.F., 2013a. Dietary lipid
732 sources as a means of changing fatty acid composition in fish: implications for
733 food fortification, in: Preedy, V.R., Srirajaskanthan, R., Patel, V.B. (Eds),
734 *Handbook of food fortification and health, from concepts to public health*
735 *applications*, volume 2. Humana Press, New York, pp. 41 – 54.

736 Pérez-Sánchez, J., Borrel, M., Bermejo-Nogales, A., Benedito-Palos, L., Saera-Vila, A.,
737 Calduch-Giner, J.A., Kaushik, S., 2013b. Dietary oils mediate cortisol kinetics
738 and the hepatic expression profile of stress responsive genes in juveniles of
739 gilthead sea bream (*Sparus aurata*) exposed to crowding stress. *Comparative*
740 *Biochemistry and Physiology D* 8, 123-130.

741 Pérez-Sánchez, J., Estensoro, I., Redondo, M.J., Calduch-Giner, J.A., Kaushik, S., Sitjà-
742 Bobadilla, A., 2013c. Mucins as diagnostic and prognostic biomarkers in a fish-
743 parasite model: transcriptional and functional analysis. *PLOS One* 8, e65457.

744 Pérez-Sánchez, J., Terova, G., Simó-Mirabet, P., Rimoldi, S., Folkedal, O., Calduch-
745 Giner, J.A., Olsen, R.E., Sitjà-Bobadilla, A., 2017. Skin mucus of gilthead sea
746 bream (*Sparus aurata* L.). Protein mapping and regulation in chronically stressed
747 fish. *Frontiers in Physiology* 8, 34.

748 Pfeiffer, C., Bauer, T., Surek, B., Schomig, E., Grundemann, D., 2011. Cyanobacteria
749 produce high levels of ergothioneine. *Food Chem.* 129, 1766-1769.

750 Piazzon, M.C., Calduch-Giner, J.A., Fouz, B., Estensoro, I., Simó-Mirabet, P., Puyalto,
751 M., Karalazos, V., Palenzuela, O., Sitjà-Bobadilla, A., Pérez-Sánchez, J., 2017.
752 Under control: how a dietary additive can restore the gut microbiome and
753 proteomic profile, and improve disease resilience in a marine teleostean fish fed
754 vegetable diets. *Microbiome* 5, 164.

755 Piazzon, C., Galindo-Villegas, J., Pereiro, P., Estensoro, I., Calduch-Giner, J.A.,
756 Gómez-Casado, E., Novoa, B., Mulero, V., Sitjà-Bobadilla, A., Pérez-Sánchez, J.,
757 2016. Differential modulation of IgT and IgM upon parasitic, bacterial, viral and
758 dietary challenges in a perciform fish. *Frontiers in Immunology* 7, 637.

759 Popowich, A., Zhang, Q., Le, X.C., 2016. Arsenobetaine: the ongoing mystery. *Nat.*
760 *Sci. Rev.* 3, 451-458.

761 Portolés, T., Ibáñez, M., Garlito, B., Nácher-Mestre, J., Karalazos, V., Silva, J., Serrano,
762 R., Pérez-Sánchez, J., Hernández, F., Berntssen, M.H.G., 2017. Comprehensive
763 strategy for pesticide residue analysis through the production cycle of gilthead sea
764 bream and Atlantic salmon. *Chemosphere* 179, 242-253.

765 Raux, E., Schubert, H.L., Warren, M.J., 2000. Biosynthesis of cobalamin (vitamin B₁₂):
766 A bacterial conundrum. *Cell. Mol. Life Sci.* 57, 1880-1893.

767 Rimoldi, S., Finzi, G., Ceccotti, C., Girardello, R., Grimaldi, A., Ascione, C., Terova,
768 G., 2016. Butyrate and taurine exert a mitigating effect on the inflamed distal
769 intestine of European sea bass fed with a high percentage of soybean meal, *Fish.*
770 *Aquat. Sci.* 19, 40.

771 Robles, R., Lozano, A.B., Sevilla, A., Marquez, L., Nuez-Ortín, W., Moyano, F.J.,
772 2013. Effect of partially protected butyrate used as feed additive on growth and
773 intestinal metabolism in sea bream (*Sparus aurata*). *Fish Physiology and*
774 *Biochemistry* 39, 1567-1580.

775 Said, H.M., Mohammed, Z.M., 2006. Intestinal absorption of water-soluble vitamins: an
776 update. *Curr. Opin. Gastroenterol.* 22, 140-146.

777 Salze, G.P., Davis, D.A., 2015. Taurine: a critical nutrient for future fish feeds.
778 *Aquaculture* 437, 215-229.

779 Sasso, O., Pontis, S., Armirotti, A., Cardinali, G., Kovacs, D., Migliore, M., Summa,
780 M., Moreno-Sanz, G., Picardo, M., Piomelli, D., 2016. Endogenous N-acyl
781 taurines regulate skin wound healing, *Proc. Natl. Acad. Sci. U. S. A.* 113, E4397-
782 E4406.

783 Simó-Mirabet, P., Felip Edo, A., Estensoro, I., Martos-Sitcha, J.A., De las Heras, V.,
784 Calduch-Giner, J., Puyalto, M., Karalazos, V., Sitjà-Bobadilla, A., Pérez-Sánchez,
785 J. 2018. Impact of low fish meal and fish oil diets on the performance, sex steroid
786 profile and male-female sex reversal of gilthead sea bream (*Sparus aurata*) over a
787 three-year production cycle. *Aquaculture* 490, 64-74.

788 Simó-Mirabet, P., Piazzon, M.C., Calduch-Giner, J.A., Ortiz, A., Puyalto, M., Sitjà-
789 Bobadilla, A., Pérez-Sánchez, J., 2017. Sodium salt medium-chain fatty acids and
790 *Bacillus*-based probiotics strategies to improve growth and intestinal health of
791 gilthead sea bream (*Sparus aurata*). *PeerJ* 5, e4001.

792 Stabler, S.P., Allen R.H., 2004. Vitamin B12 deficiency as a world-wide problem.
793 *Annu. Rev. Nutr.* 24, 299-326.

794 Stiboller, M., Raber, G., Francesconi, K.A., 2015. Simultaneous determination of
795 glycine betaine and arsenobetaine in biological samples by HPLC/ICPMS/ESMS
796 and the application to some marine and freshwater fish samples. *Microchem. J.*
797 122, 172-175.

798 Tacon, A.G.J., Metian, M., 2008. Global overview on the use of fish meal and fish oil in
799 industrially compounded aquafeeds: Trends and future prospects. *Aquaculture*
800 285, 146-158.

801 Tai, S.S.-C., Bedner, M., Phinney, K.W., 2010. Development of a candidate reference
802 measurement procedure for the determination of 25-hydroxyvitamin D₃ and 25-
803 hydroxyvitamin D₂ in human serum using isotope-dilution liquid
804 chromatography–tandem mass spectrometry. *Analytical Chemistry* 82, 1942-
805 1948.

806 Tocher, D.R., 2015. Omega-3 long-chain polyunsaturated fatty acids and aquaculture in
807 perspective. *Aquaculture* 449, 94-107. Turchini, G.M., Hermon, K.M., Francis,
808 D.S., 2018. Fatty acids and beyond: Fillet nutritional characterisation of rainbow
809 trout (*Oncorhynchus mykiss*) fed different dietary oil sources. *Aquaculture* 491,
810 391-397.

811 Une, M., Goto, T., Kihira, K., Kuramoto, T., Hagiwara, K., Nakajima, T., Hoshita, T.,
812 1991. Isolation and identification of bile salts conjugated with cysteinolic acid
813 from bile of the red seabream, *Pagrosomus major*. *J. Lipid Res.* 32, 1619-1623.

814 Venter, L., Jansen van Rensburg, P., Loots, D.T., Vosloo, A., Lindeque, J.Z., 2016.
815 Untargeted metabolite profiling of abalone using gas chromatography mass
816 spectrometry. *Food Anal. Methods* 9, 1254-1261.

817 Watanabe, F., Nakano, Y., Tachikake, N., Saido, H., Tamura, Y., Yamanaka, H., 1991.
818 Vitamin B-12 deficiency increases the specific activities of rat liver NADH- and
819 NADPH-linked aquacobalamin reductase isozymes involved in coenzyme
820 synthesis. *J. Nutr.* 121, 1948-1954.

821 Wiklund, S., Johansson, E., Sjöström, L., Mellerowicz, E.J., Edlund, U., Shockcor, J.P.,
822 Gottfries, J., Moritz, T., Trygg J., 2008. Visualization of GC/TOF-MS-based
823 metabolomics data for identification of biochemically interesting compounds
824 using OPLS class models. *Analytical Chemistry* 80, 115-122.

825 Wishart, D.S., Jewison, T., Guo, A.C., Wilson, M., Knox, C., Liu, Y., Djoumbou, Y.,
826 Mandal, R., Aziat, F., Dong, E., Bouatra, S., Sinelnikov, I., Arndt, D., Xia, J., Liu,
827 P., Yallou, F., Bjorndahl, T., Perez-Pineiro, R., Eisner, R., Allen, F., Neveu, V.,
828 Greiner, R., Scalbert, A., 2013 HMDB 3.0--The Human Metabolome Database in
829 2013. *Nucleic Acids Res.* 41, D801-D807.

830 Wold, S., Sjöström, M., Eriksson, L., 2001. PLS-regression: a basic tool of
831 chemometrics. *Chemometrics Intelligent Lab. Syst.* 58, 109–130. Yancey, P.H.,
832 2005. Organic osmolytes as compatible, metabolic and counteracting
833 cytoprotectants in high osmolarity and other stresses. *J. Exp. Biol.* 208, 2819-
834 2830.

Figure captions

Fig. 1. PLS-DA score plot of acquired data of D1 group individuals (black), D2 (red) and D3/4 (green for D3, blue for D4). Insert is a screen plot of the principal component analysis, showing eigenvalues (blue bars) and cumulative variability explained (orange points) against the number of the principal component.

Fig. 2. Correlation plot of hercynine integrated area in feeds (X-axis) and individual serum samples (Y-axis).

Fig. 3. Integrative profile of differential compounds between D2 and D3/4 compared to control D1 group. Bars show for each dietary group and biological process the number of significantly different ($P < 0.5$, ANOVA followed by Benjamini-Hochberg multiple testing correction) abundant compounds. Colors in each bar indicate the level of change (as % of D1) as indicated in the inbox.

850 **Table 1.** Ingredients and chemical composition of experimental diets.

Ingredient (%)	Diet			
	D1	D2	D3	D4
Fish meal	23.0	3.0	3.0	3.0
Fish hydrolysate (CPSP)	2.0	2.0	2.0	2.0
Soya protein	16.0	25.0	25.0	25.0
Corn gluten	15.0	25.0	25.0	25.0
Wheat gluten	4.0	7.3	7.3	7.3
Rapeseed cake	12.0	9.7	9.9	9.9
Wheat	11.08	6.80	6.64	6.24
Fish oil	15.60	6.56	2.50	2.50
Rapeseed oil	0.0	4.4	6.5	6.5
Palm oil	0.0	4.4	6.5	6.5
Monocalcium phosphate	0.303	2.097	2.097	2.097
Histidine	0.136	0.136	0.136	0.136
Mineral Vitamin mix ^a	0.5	0.5	0.5	0.5
Cholesterol	0.113	0.113	0.113	0.113
Amino-acid and micronutrient mix ^b	0.20	2.92	2.74	2.74
Antioxidants	0.045	0.045	0.045	0.045
Yttrium	0.03	0.03	0.03	0.03
Butyrate (BP-70)	0.0	0.0	0.0	0.4
<i>Proximate composition</i>				
Dry matter (DM, %)	91.65	91.79	91.80	92.34
Crude protein (%DM)	45.48	46.73	46.12	46.03
Crude fat (% DM)	19.80	19.56	20.13	19.40
EPA + DHA (% DM)	2.90	1.38	0.67	0.63

851 ^a Supplied the following (g/kg mix, except as noted): calcium 689, sodium 108, iron 3,
852 manganese 1, zinc 1, cobalt 2 mg, iodine 2 mg, selenium 20 mg, molybdenum 32 mg, retinyl
853 acetate 1, DL-cholecalciferol 2.6, DL- α tocopheryl acetate 28, menadione sodium bisulphite 2,
854 ascorbic acid 16, thiamin 0.6, riboflavin 1.7, pyridoxine 1.2, vitamin B₁₂ 50 mg, nicotinic acid 5,
855 pantothenic acid 3.6, folic acid 0.6, and biotin 50 mg.

856 ^b Contains methionine, lysine, choline, and lecithin.

857 **Table 2.** Biometry of sampled gilthead sea bream fed experimental diets. Values are the
858 mean \pm SEM (n= 10).

	D1	D2	D3	D4	P-value (ANOVA)
Body weight (g)	611.95 \pm 24.2	587.40 \pm 25.8	580.8 \pm 10.7	577.6 \pm 21.0	0.679
Liver weight (g)	7.33 \pm 0.33	7.42 \pm 0.64	8.06 \pm 0.38	7.38 \pm 0.38	0.855
Mesenteric fat (g)	13.80 \pm 2.18	11.89 \pm 2.16	10.61 \pm 1.41	10.38 \pm 1.50	0.546
HSI (%) ¹	1.20 \pm 0.05	1.27 \pm 0.06	1.39 \pm 0.06	1.28 \pm 0.06	0.124
MSI (%) ²	2.20 \pm 0.31	2.19 \pm 0.28	1.80 \pm 0.20	1.79 \pm 0.25	0.673

859 ¹Hepatosomatic index = (100 x liver weight) / fish weight.

860 ²Mesenteric fat index = (100 x mesenteric fat) / fish weight.

861 **Table 3.** Highlighted compounds obtained from untargeted metabolomics. Values are the mean \pm SEM (n= 8-10).

Compound name		Biological process [†]	Chromatography/ionization mode	Formula	De/protonated molecule <i>m/z</i> (mDa)	RT (min)	D2, % CTRL	D3/4, %CTRL	Corrected P-value [‡]	VIP ^{††}
1	PC(22:6/16:0)	1	RP(spec) / +	C ₄₆ H ₈₀ NPO ₈	806.5701 (+0.1)	18.86	68 \pm 5 ^b	39 \pm 7 ^c	1.63E ⁻⁰⁶	2.12
2	PC(22:6/18:0)	1	RP(spec) / +	C ₄₈ H ₈₄ NPO ₈	834.6010 (-0.3)	19.86	71 \pm 13 ^b	49 \pm 6 ^c	3.57E ⁻⁰⁶	1.48
3	PC(22:6/18:3)	1	RP(spec) / +	C ₄₈ H ₇₈ NPO ₈	828.5544 (+0.1)	17.65	94 \pm 15 ^a	49 \pm 4 ^b	1.54E ⁻⁰³	1.47
4	PC(22:6/20:4)	1	RP(spec) / +	C ₅₀ H ₈₀ NPO ₈	854.5700 (+0.1)	19.05	38 \pm 3 ^b	16 \pm 2 ^c	7.16E ⁻¹⁰	1.90
5	PC(22:6/20:5)	1	RP(spec) / +	C ₅₀ H ₇₈ NPO ₈	852.5541 (-0.2)	17.54	58 \pm 4 ^b	15 \pm 2 ^c	1.91E ⁻⁰⁹	1.33
6	PC(20:5/14:0)	1	RP(spec) / +	C ₄₂ H ₇₂ NPO ₈	750.5079 (+0.5)	23.44	32 \pm 4 ^b	13 \pm 1 ^c	1.62E ⁻¹⁵	2.03
7	PC(20:5/16:0)	1	RP(spec) / +	C ₄₄ H ₇₈ NPO ₈	780.5532 (-1.1)	18.45	82 \pm 8 ^b	30 \pm 5 ^c	4.31E ⁻¹¹	1.93
8	PC(20:5/16:1)	1	RP(spec) / +	C ₄₄ H ₇₆ NPO ₈	778.5385 (-0.2)	17.88	33 \pm 3 ^b	15 \pm 1 ^c	1.82E ⁻¹⁵	1.86
9	PC(20:5/18:0)	1	RP(spec) / +	C ₄₆ H ₈₂ NPO ₈	808.5855 (-0.1)	19.5	64 \pm 12 ^b	39 \pm 1 ^c	8.38E ⁻⁰⁷	1.96
10	PC(20:5/18:1)	1	RP(spec) / +	C ₄₆ H ₈₀ NPO ₈	806.5700 (0.0)	18.56	107 \pm 14 ^a	80 \pm 6 ^b	6.59E ⁻⁰²	1.47
11	PC(20:5/18:2)	1	RP(spec) / +	C ₄₆ H ₇₈ NPO ₈	804.5541(-0.2)	17.88	51 \pm 8 ^b	20 \pm 2 ^c	4.98E ⁻⁰⁷	1.72
12	PC(20:5/18:3)	1	RP(spec) / +	C ₄₆ H ₇₆ NPO ₈	802.5277 (-1.0)	18.43	125 \pm 18 ^b	63 \pm 12 ^c	3.84E ⁻⁰²	1.38
13	PC(20:5/20:4)	1	RP(spec) / +	C ₄₈ H ₇₆ NPO ₈	826.5381 (-0.6)	17.25	92 \pm 8 ^a	46 \pm 6 ^b	4.98E ⁻⁰⁴	1.33
14	PC(20:5/20:5)	1	RP(spec) / +	C ₄₈ H ₇₄ NPO ₈	824.5220 (-1.0)	17.28	43 \pm 4 ^b	9 \pm 1 ^c	3.26E ⁻⁰⁸	1.66
15	PC(18:2/16:0)	1	RP(spec) / +	C ₄₂ H ₈₀ NPO ₈	758.5701 (+0.1)	19.23	410 \pm 33 ^b	571 \pm 74 ^c	3.07E ⁻¹⁴	2.13
16	PC(18:2/18:0)	1	RP(spec) / +	C ₄₄ H ₈₄ NPO ₈	786.6008 (-0.5)	20.34	625 \pm 94 ^b	1689 \pm 270 ^c	1.31E ⁻⁰⁵	2.12
17	PC(18:2/18:2)	1	RP(spec) / +	C ₄₄ H ₈₀ NPO ₈	782.5711 (+1.1)	18.6	568 \pm 68 ^b	1693 \pm 271 ^c	3.16E ⁻⁰⁸	1.67
18	PC(18:1/16:0)	1	RP(spec) / +	C ₄₂ H ₈₂ NPO ₈	760.5859 (+0.3)	20.03	151 \pm 20 ^b	193 \pm 14 ^c	2.44E ⁻⁰³	1.31
19	PC(18:1/18:0)	1	RP(spec) / +	C ₄₄ H ₈₆ NPO ₈	788.6191 (+2.2)	21.22	204 \pm 33 ^b	316 \pm 35 ^c	6.10E ⁻⁰³	1.89

20	PC(18:1/18:1)	1	RP(spec) / +	C ₄₄ H ₈₄ NPO ₈	786.6012 (-0.1)	20.31	483 ± 68 ^b	761 ± 68 ^c	1.10E ⁻⁰⁶	2.12
21	PC(18:1/18:2)	1	RP(spec) / +	C ₄₄ H ₈₂ NPO ₈	784.5858 (+0.2)	19.37	766 ± 130 ^b	1581 ± 285 ^c	6.29E ⁻⁰⁹	2.12
22	PC(18:1/18:3)	1	RP(spec) / +	C ₄₄ H ₈₀ NPO ₈	782.5712 (+1.2)	18.6	488 ± 93 ^b	1419 ± 199 ^c	6.43E ⁻⁰⁸	1.67
23	PC(16:0/18:0)	1	RP(spec) / +	C ₄₂ H ₈₄ NPO ₈	762.6013 (0.0)	20.03	517 ± 36 ^b	1080 ± 119 ^c	1.74E ⁻¹⁸	2.16
24	PC(16:0/18:3)	1	RP(spec) / +	C ₄₂ H ₇₈ NPO ₈	756.5545 (+0.2)	18.5	532 ± 85 ^b	1317 ± 105 ^c	3.84E ⁻⁰⁸	1.96
25	LysoPC(22:6)	1	RP(spec) / +	C ₃₀ H ₅₀ NPO ₇	568.3405 (+0.2)	9.81	77 ± 15 ^b	60 ± 8 ^b	1.21E ⁻⁰⁵	1.55
26	LysoPC(22:5)	1	RP(spec) / +	C ₃₀ H ₅₂ NPO ₇	570.3566 (+0.6)	9.11	143 ± 20 ^b	157 ± 25 ^b	7.17E ⁻⁰³	1.37
27	LysoPC(20:5)	1	RP(spec) / +	C ₂₈ H ₄₈ NPO ₇	542.3242 (+0.5)	8.58	88 ± 9 ^b	87 ± 7 ^b	4.21E ⁻⁰²	1.59
28	LysoPC(20:4)	1	RP(spec) / +	C ₂₈ H ₅₀ NPO ₇	544.3386 (-1.7)	8.12	50 ± 10 ^b	32 ± 4 ^b	2.93E ⁻⁰⁸	133
29	LysoPC(20:2)	1	RP(spec) / +	C ₂₈ H ₅₄ NPO ₇	548.3714 (-0.2)	7.88	294 ± 50 ^b	468 ± 47 ^c	1.62E ⁻¹⁵	1.61
30	LysoPC(18:3)	1	RP(spec) / +	C ₂₆ H ₄₈ NPO ₇	518.3248 (+0.1)	10.82	212 ± 17 ^b	328 ± 66 ^c	1.69E ⁻¹⁴	1.54
31	LysoPC(18:2)	1	RP(spec) / +	C ₂₆ H ₅₀ NPO ₇	520.3403 (0.0)	9.00	237 ± 28 ^b	398 ± 40 ^c	5.03E ⁻¹¹	1.71
32	LysoPC(18:1)	1	RP(spec) / +	C ₂₆ H ₅₂ NPO ₇	522.3557 (-0.3)	8.41	138 ± 18 ^b	195 ± 29 ^c	8.14E ⁻⁰⁷	1.73
33	LysoPC(18:0)	1	RP(spec) / +	C ₂₆ H ₅₄ NPO ₇	524.3704 (-1.2)	7.55	113 ± 10 ^b	170 ± 19 ^c	3.78E ⁻⁰⁴	1.43
34	LysoPC(16:0)	1	RP(spec) / +	C ₂₄ H ₅₀ NPO ₇	496.3402 (-0.1)	10.82	145 ± 26 ^b	323 ± 42 ^c	3.33E ⁻⁰⁵	1.38
35	FFA(22:6)	2	RP / -	C ₂₂ H ₃₂ O ₂	327.2316 (-0.8)	15.18	85 ± 10 ^a	67 ± 13 ^b	3.25E ⁻⁰³	1.31
36	FFA(20:5)	2	RP / -	C ₂₀ H ₃₀ O ₂	301.2167 (-0.1)	15.17	96 ± 17 ^a	77 ± 12 ^b	4.00E ⁻⁰³	1.55
37	FFA(20:4)	2	RP / -	C ₂₀ H ₃₂ O ₂	303.2316 (-0.8)	15.86	92 ± 12 ^a	78 ± 5 ^b	9.00E ⁻⁰³	1.77
38	FFA(18:4)	2	RP / -	C ₁₈ H ₂₈ O ₂	275.2004 (-0.7)	14.98	51 ± 7 ^b	27 ± 5 ^c	7.51E ⁻¹⁶	1.95
39	FFA(18:2)	2	RP / -	C ₁₈ H ₃₂ O ₂	279.2316 (-0.8)	15.91	202 ± 38 ^b	295 ± 21 ^c	6.03E ⁻⁰⁹	1.95
40	FFA(18:1)	2	RP / -	C ₁₈ H ₃₄ O ₂	281.2472 (-0.9)	15.66	160 ± 18 ^b	308 ± 34 ^c	1.32E ⁻⁰⁴	1.45
41	FFA(16:1)	2	RP / -	C ₁₆ H ₃₀ O ₂	253.2161 (-0.7)	16.43	103 ± 19 ^a	177 ± 25 ^c	3.00E ⁻⁰³	1.35

42	FFA(16:0)	2	RP / -	C ₁₆ H ₃₂ O ₂	255.2316 (-0.8)	16.43	111 ± 18 ^a	136 ± 23 ^b	1.05E ⁻⁰²	1.40
43	(9-methyl-d19:3) sphingosine	3	RP / +	C ₁₉ H ₃₇ NO ₂	312.2899 (-0.4)	12.32	23 ± 3 ^b	16 ± 2 ^c	1.71E ⁻¹¹	2.11
44	(D14:2)sphingosine	3	RP / +	C ₁₄ H ₂₇ NO ₂	242.2118 (-0.2)	9.17	878 ± 123 ^b	148 ± 27 ^c	1.98E ⁻⁰⁴	2.06
45	N-Heptadecenoyl taurine	4	RP / -	C ₁₉ H ₃₇ NSO ₄	374.2355 (-1.0)	15.08	52 ± 10 ^b	22 ± 2 ^c	6.04E ⁻¹⁴	1.90
46	N-Palmitoleoyl taurine	4	RP / -	C ₁₈ H ₃₅ NSO ₄	360.2209 (0.0)	14.58	47 ± 4 ^b	22 ± 3 ^c	2.87E ⁻¹²	1.95
47	Cytidine	5	HI / +	C ₉ H ₁₃ N ₃ O ₅	244.0941 (+0.8)	4.37	235 ± 28 ^b	130 ± 21 ^c	1.34E ⁻⁰²	1.39
48	Cytosine	5	HI / +	C ₄ H ₅ N ₃ O	112.0502 (-0.9)	4.35	200 ± 26 ^b	120 ± 8 ^c	1.07E ⁻⁰²	1.78
49	Deoxycytidine	5	HI / +	C ₉ H ₁₃ N ₃ O ₄	228.0951 (-3.3)	3.48	653 ± 59 ^b	190 ± 27 ^c	5.62E ⁻⁰³	2.27
50	Methylcytosine	5	HI / +	C ₅ H ₇ N ₃ O	126.0645 (-2.2)	4.29	734 ± 103 ^b	120 ± 24 ^c	3.87E ⁻⁰⁵	2.13
51	Cysteinolic acid	6,10	HI / -	C ₃ H ₉ NSO ₄	154.0169 (-0.5)	4.05	18 ± 3 ^b	10 ± 2 ^b	6.51E ⁻¹⁵	2.33
52	Tauropine	7,10	HI / -	C ₅ H ₁₁ NSO ₅	196.0286 (+0.6)	2.28	28 ± 4 ^b	35 ± 7 ^b	1.74E ⁻¹⁰	2.32
53	TMAO	7,10	HI / +	C ₃ H ₉ NO	76.0760 (-0.2)	5.87	53 ± 10 ^b	49 ± 9 ^b	2.69E ⁻⁰³	1.52
54	Arsenobetaine	8,10	HI / +	C ₅ H ₁₁ AsO ₂	179.0040 (-1.3)	5.78	50 ± 6 ^b	52 ± 7 ^b	5.00E ⁻⁰⁵	1.97
55	Hercynine	9,10	HI / +	C ₉ H ₁₅ N ₃ O ₂	198.1235 (-0.8)	5.75	737 ± 74 ^b	182 ± 15 ^c	3.00E ⁻⁰⁶	2.55

1, Phospholipid metabolism; 2, Fatty acid metabolism; 3, Sphingolipid metabolism; 4, N-acyl amino acid metabolism; 5, Pyrimidine metabolism; 6, Bile acid metabolism/algae amino acid; 7, Anaerobic microbial metabolism; 8, Arsenic metabolism; 9, Fungi metabolism; 10, Exogenous compounds.

[‡]ANOVA followed by Benjamini-Hochberg multiple testing correction. ^{††}Variable importance in projection measurements in PLS-DA.

865 **Table 4.** Vitamin and vitamin-related compounds obtained from refined targeted approach. Values are the mean \pm SEM (n= 8-10).
866

Vitamin/vitamin-related compounds		Chromatography/ ionization mode	Formula	De/protonated molecule <i>m/z</i> (error mDa)	RT (min)	(%) CTRL D2 [†]	(%) CTRL D3/4 [†]	P-value (ANOVA)
A	Retinol phosphate	RP/+	C ₂₀ H ₃₁ O ₄ P	367.2015 (-2.3)	15.75	140 \pm 60 ^a	121 \pm 63 ^a	4.45E-01
B ₁	Thiamin	HI/+	C ₁₂ H ₁₆ N ₄ OS	265.1118 (-0.5)	5.68	120 \pm 18 ^a	78 \pm 23 ^a	2.29E-01
B ₂	Riboflavin	RP/-	C ₁₇ H ₂₀ N ₄ O ₆	375.1299 (-0.6)	4.44	144 \pm 67 ^a	364 \pm 132 ^b	1.56E-03
B ₅	Pantothenic acid	RP/+	C ₉ H ₁₇ NO ₅	220.1183 (-0.2)	2.04	120 \pm 17 ^a	146 \pm 25 ^b	1.98E-02
B ₆	Pyridoxine	RP/+	C ₈ H ₁₁ NO ₃	170.0829 (+1.2)	1.72	96 \pm 24 ^a	104 \pm 21 ^a	5.96E-01
B ₇	Biotin	RP/+	C ₁₀ H ₁₆ N ₂ O ₃ S	245.0955 (-0.5)	5.36	107 \pm 19 ^a	120 \pm 21 ^a	3.68E-01
B ₁₂	Mehtylmalonic acid (MMA)	RP/-	C ₄ H ₆ O ₄	117.0190 (+0.2)	1.22	195 \pm 45 ^b	276 \pm 35 ^c	3.27E-03
C	Dehydroascorbic acid	HI/-	C ₆ H ₆ O ₆	173.0085 (-0.1)	1.12	95 \pm 17 ^a	122 \pm 17 ^a	1.03E-01
D ₃	25-hydroxyvitamin D ₃	RP/+	C ₂₇ H ₄₄ O ₂	401.3412 (-0.8)	13.65	102 \pm 39 ^a	93 \pm 26 ^a	3.59E-01
E	α -Carboxyethylhydroxychroman	RP/-	C ₁₆ H ₂₂ O ₄	277.1441 (+0.1)	14.10	106 \pm 17 ^a	109 \pm 15 ^a	5.64E-01
K ₂	Menaquinone	RP/+	C ₄₁ H ₅₆ O ₂	581.4360 (+0.1)	16.80	72 \pm 45 ^a	140 \pm 54 ^a	1.07E-01

867 [†] Percentage of integrated area for the selected compound as a percentage in fish fed control diet (D1). Compounds with statistical significant differences (P< 0.05) against
868 control fish are in bold.
869